

SRI International

N93-27000

Unclass

G3/25 0163207

Final Report • April 1993

REPORT ON CHEMICAL ANALYSES OF PROVIDED SAMPLES

Christopher H. Becker
Molecular Physics Laboratory

SRI Project 3557
Contract No. 1-1EH-46755

MP 93-084

Prepared for:

NASA Marshall Space Flight Center
Huntsville, AL 35812

Attn: Roger Linton
Mail Stop EH15

Approved:

Donald J. Eckstrom, Director
Molecular Physics Laboratory

1N-25-CR
163 207
P.19

(NASA-CR-193207) CHEMICAL ANALYSES
OF PROVIDED SAMPLES Final Report
(SRI International Corp.) 19 p

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small>				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE April 1993	3. REPORT TYPE AND DATES COVERED Final Report		
4. TITLE AND SUBTITLE REPORT ON CHEMICAL ANALYSES OF PROVIDED SAMPLES		5. FUNDING NUMBERS		
6. AUTHOR(S) Christopher H. Becker				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) SRI International 333 Ravenswood Avenue Menlo Park, CA 94025		8. PERFORMING ORGANIZATION REPORT NUMBER PYU 3557 MP 93-084		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) NASA Marshall Space Flight Center Huntsville, AL 35812		10. SPONSORING/MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) A batch of four samples were received and chemical analysis was performed of the surface and near surface regions of the samples by the surface analysis by laser ionization (SALI) method. The samples included four one-inch diameter optics labeled windows # PR14 and PR17 and MgF2 mirrors 9-93 PPCC exp. and control DMES 26-92. The analyses emphasized surface contamination or modification. In these studies, pulsed desorption by 355 nm laser light and single-photon ionization (SPI) above the sample by coherent 118 nm radiation (at ~5 x 10 ⁵ W/cm ²) were used, emphasizing organic analysis. For the two windows with an apparent yellowish contaminant film, higher desorption laser power was needed to provide substantial signals, indicating a less volatile contamination than for the two mirrors. Window PR14 and the 9-93 mirror showed more hydrocarbon components than the other two samples. The mass spectra, which show considerable complexity, are discussed in terms of various potential chemical assignments.				
14. SUBJECT TERMS EDEF, surface analysis, surface contamination, space environmental effects			15. NUMBER OF PAGES 17	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL	

A batch of four samples was received consisting of one inch diameter optics labeled windows # PR14 and PR17, and MgF2 mirrors 9-93 PPC exp. and control DMES 26-92. Chemical analyses of the surfaces were performed by the surface analysis by laser ionization (SALI) method. The analyses emphasize surface contamination. SALI uses nonselective photoionization of sputtered or desorbed atoms and molecules above but close (~ 1 mm) to the surface, followed by time-of-flight (TOF) mass spectrometry. In these studies, laser-induced desorption by 5 ns pulse-width 355 nm light ($10\text{--}100$ mJ/cm²) and single-photon ionization (SPI) by coherent 118 nm radiation (at $\sim 5 \times 10^5$ W/cm²) were used. SPI was chosen primarily for its ability to obtain molecular information, whereas multiphoton ionization (not used in the present studies) is intended primarily for elemental and small molecule information. The choice of laser desorption was made after initial studies with pulsed Ar⁺ sputtering; it was apparent that considerably more dynamic range and higher mass range were available with laser desorption; also charging of the optics was occurring with the pulsed Ar⁺ beam.


While SPI by 118 nm (10.5 eV) light is considered a generally "soft" (nonfragmenting) form of radiation, stimulated desorption can cause fragmentation and also produce internally hot molecules which photofragment relatively easily compared to lower temperature sources of molecules. Typically, the low mass regions of the mass spectra contain a good deal of molecular fragment information.

Some other comments on the mass spectra are appropriate. The signals from the microchannel plate detector were recorded in analog fashion by a 100 MHz transient digitizer; thus the voltage signals are given as "relative intensity" and not as ion counts. For the two windows with apparent yellowish contaminant films (PR14 and 17), higher desorption laser intensities were needed to provide substantial signals, indicating a less volatile contamination than for the two mirrors (or possibly poorer absorptivity of 355 nm light). For sample PR17, and a little bit for the control mirror (DMES 26-92), direct ion signals (intense and broad mass peaks on a different mass scale--different time zero) are seen in the low mass region.


Some of the mass assignments have been marked on the figures, especially for the major peaks. Considerable effort can be invoked for mass interpretation. Some possible brief general interpretations now follow. Window PR14 and the 9-93 MgF2 mirror show considerably more hydrocarbon fragments at low masses and thus likely contain much more hydrocarbon components than the other two samples.

For PR17 the m/z 105 and 207 peaks can be associated with Si and O ($\text{Si}_2\text{O}_3\text{H}$ for 105 and $\text{Si}_3\text{O}_3(\text{CH}_3)_5$ for 207), although there are other interpretations such as $\text{C}_6\text{H}_5\text{-CO}$ for m/z 105. Also it would be unusual to observe a strong 207 and not a strong 73 or 147 for a silicone. Polydimethylsilicone was run as a standard for comparison under these conditions; strong peaks at m/z 73, 147, 207, 221, and 281 were observed.


For the two MgF2 mirrors, there are numerous major mass peaks in common such as 43, 57, 97, 112, 149, 167, 185 and 221. Note that 221 can possibly be assigned to a silicone, but the other peaks are not commonly associated with silicones except m/z 129 in the 9-93 mirror which can be rationalized as a $\text{Si}_2\text{C}_5\text{H}_{13}$ structure. Masses 97, 112, 149, 167, and 185 can be assigned to $\text{C}_x\text{F}_y\text{O}_z$ structures. As to the higher mass assignments for the control mirror, it is not clear (especially with no detailed knowledge of sample history) what structures are associated with these masses; however, these peaks will be very characteristic of whatever compound(s) is present.



MgF2 MIRROR CONTROL, DMES 26-92, 355+118 NM, AP1417




PR14 WINDOW, 355 + 118 NM, AP1404



AP1404


PR14

16-APR-93 18:42:25




AP1404 PR14


16-APR-93 18:44:54




AP1404 PR14 16-APR-93 18:46:50




PR17 WINDOW, 355 + 118 NM, AP1410




PR17 WINDOW, 355 + 118 NM, AP1410




PR17 WINDOW, 355 + 118 NM, AP1410




MgF2 MIRROR 9-93 PPPC exp., 355+118 NM, AP1414




MgF2 MIRROR 9-93 PPPC exp., 355+118 NM, AP1414




MgF2 MIRROR 9-93 PPPC exp., 355+118 NM, AP1414




MgF2 MIRROR 9-93 PPPC exp., 355+118 NM, AP1414



MgF2 MIRROR CONTROL, DMES 26-92, 355+118 NM, AP1417



MgF2 MIRROR CONTROL, DMES 26-92, 355+118 NM, AP1417



MgF2 MIRROR CONTROL, DMES 26-92, 355+118 NM, AP1417

SRI International

93-27000

Final Report • April 1993

REPORT ON CHEMICAL ANALYSES OF PROVIDED SAMPLES

Christopher H. Becker
Molecular Physics Laboratory

SRI Project 3557
Contract No. 1-1EH-46755

MP 93-084

Prepared for:

NASA Marshall Space Flight Center
Huntsville, AL 35812

Attn: Roger Linton
Mail Stop EH15

Approved:

Donald J. Eckstrom, Director
Molecular Physics Laboratory

A batch of four samples was received consisting of one inch diameter optics labeled windows # PR14 and PR17, and MgF2 mirrors 9-93 PPC exp. and control DMES 26-92. Chemical analyses of the surfaces were performed by the surface analysis by laser ionization (SALI) method. The analyses emphasize surface contamination. SALI uses nonselective photoionization of sputtered or desorbed atoms and molecules above but close (~ 1 mm) to the surface, followed by time-of-flight (TOF) mass spectrometry. In these studies, laser-induced desorption by 5 ns pulse-width 355 nm light (10 - 100 mJ/cm²) and single-photon ionization (SPI) by coherent 118 nm radiation (at $\sim 5 \times 10^5$ W/cm²) were used. SPI was chosen primarily for its ability to obtain molecular information, whereas multiphoton ionization (not used in the present studies) is intended primarily for elemental and small molecule information. The choice of laser desorption was made after initial studies with pulsed Ar⁺ sputtering; it was apparent that considerably more dynamic range and higher mass range were available with laser desorption; also charging of the optics was occurring with the pulsed Ar⁺ beam.


While SPI by 118 nm (10.5 eV) light is considered a generally "soft" (nonfragmenting) form of radiation, stimulated desorption can cause fragmentation and also produce internally hot molecules which photofragment relatively easily compared to lower temperature sources of molecules. Typically, the low mass regions of the mass spectra contain a good deal of molecular fragment information.


Some other comments on the mass spectra are appropriate. The signals from the microchannel plate detector were recorded in analog fashion by a 100 MHz transient digitizer; thus the voltage signals are given as "relative intensity" and not as ion counts. For the two windows with apparent yellowish contaminant films (PR14 and 17), higher desorption laser intensities were needed to provide substantial signals, indicating a less volatile contamination than for the two mirrors (or possibly poorer absorptivity of 355 nm light). For sample PR17, and a little bit for the control mirror (DMES 26-92), direct ion signals (intense and broad mass peaks on a different mass scale--different time zero) are seen in the low mass region.

Some of the mass assignments have been marked on the figures, especially for the major peaks. Considerable effort can be invoked for mass interpretation. Some possible brief general interpretations now follow. Window PR14 and the 9-93 MgF2 mirror show considerably more hydrocarbon fragments at low masses and thus likely contain much more hydrocarbon components than the other two samples.


For PR17 the m/z 105 and 207 peaks can be associated with Si and O ($\text{Si}_2\text{O}_3\text{H}$ for 105 and $\text{Si}_3\text{O}_3(\text{CH}_3)_5$ for 207), although there are other interpretations such as $\text{C}_6\text{H}_5\text{-CO}$ for m/z 105. Also it would be unusual to observe a strong 207 and not a strong 73 or 147 for a silicone. Polydimethylsilicone was run as a standard for comparison under these conditions; strong peaks at m/z 73, 147, 207, 221, and 281 were observed.

For the two MgF2 mirrors, there are numerous major mass peaks in common such as 43, 57, 97, 112, 149, 167, 185 and 221. Note that 221 can possibly be assigned to a silicone, but the other peaks are not commonly associated with silicones except m/z 129 in the 9-93 mirror which can be rationalized as a $\text{Si}_2\text{C}_5\text{H}_{13}$ structure. Masses 97, 112, 149, 167, and 185 can be assigned to $\text{C}_x\text{F}_y\text{O}_z$ structures. As to the higher mass assignments for the control mirror, it is not clear (especially with no detailed knowledge of sample history) what structures are associated with these masses; however, these peaks will be very characteristic of whatever compound(s) is present.






PR14 WINDOW, 355 + 118 NM, AP1404



AP1404


PR14

16-APR-93 18:42:25




AP1404 PR14

16-APR-93 18:44:54





AP1404 PR14


16-APR-93 18:46:50




PR17 WINDOW, 355 + 118 NM, AP1410









MgF2 MIRROR 9-93 PPC exp., 355+118 NM, AP1414




MgF2 MIRROR 9-93 PPC exp., 355+118 NM, AP1414




MGF2 MIRROR 9-93 PPC exp., 355+118 NM, AP1414




MgF2 MIRROR 9-93 PPPC exp., 355+118 NM, AP1414



MgF2 MIRROR CONTROL, DMES 26-92, 355+118 NM, AP1417



MgF2 MIRROR CONTROL, DMES 26-92, 355+118 NM, AP1417



MgF2 MIRROR CONTROL, DMES 26-92, 355+118 NM, AP1417

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503</small>				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE April 1993	3. REPORT TYPE AND DATES COVERED Final Report		
4. TITLE AND SUBTITLE REPORT ON CHEMICAL ANALYSES OF PROVIDED SAMPLES		5. FUNDING NUMBERS P.O. H-07967D		
6. AUTHOR(S) Christopher H. Becker				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) SRI International 333 Ravenswood Avenue Menlo Park, CA 94025		8. PERFORMING ORGANIZATION REPORT NUMBER PYU 3557 MP 93-084		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) NASA Marshall Space Flight Center Huntsville, AL 35812		10. SPONSORING/MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words) A batch of four samples were received and chemical analysis was performed of the surface and near surface regions of the samples by the surface analysis by laser ionization (SALI) method. The samples included four one-inch diameter optics labeled windows # PR14 and PR17 and MgF2 mirrors 9-93 PPCC exp. and control DMES 26-92. The analyses emphasized surface contamination or modification. In these studies, pulsed desorption by 355 nm laser light and single-photon ionization (SPI) above the sample by coherent 118 nm radiation (at ~5 x 10 ⁵ W/cm ²) were used, emphasizing organic analysis. For the two windows with an apparent yellowish contaminant film, higher desorption laser power was needed to provide substantial signals, indicating a less volatile contamination than for the two mirrors. Window PR14 and the 9-93 mirror showed more hydrocarbon components than the other two samples. The mass spectra, which show considerable complexity, are discussed in terms of various potential chemical assignments.				
14. SUBJECT TERMS EDEF, surface analysis, surface contamination, space environmental effects		15. NUMBER OF PAGES 17		
		16. PRICE CODE		
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL	